

キイトルーダ+パクリタキセル

適 応 疾 患：

PD-L1 陽性のホルモン受容体陰性かつ HER2 陰性 (triple-negative) の治癒切除不能又は転移・再発乳癌

1 コースの期間：12 週間 上限コース回数：なし

Rp.	薬剤	投与量	投与経路	投与速度、時間
1	生食	250mL	点滴静注	メインルート
2	レスタミンコーワ錠	50mg	内服	パクリタキセル投与30分前までに
3	デキサート注 ファモチジン注	8.25mg 20mg	静注	パクリタキセル投与約30分前に緩徐に静注
4	キイトルーダ 生食	200mg/body 100mL	点滴静注	30分
5	生食	50mL	点滴静注	30分
6	パクリタキセル 生食	90mg/m ² 250mL	点滴静注	60分

	day																											
Rp.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
1	↓							↓							↓							↓						
2	↓							↓							↓													
3	↓							↓							↓													
4	↓																					↓						
5								↓							↓													
6	↓							↓							↓													

	day																											
Rp.	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56
1	↓							↓							↓													
2	↓							↓							↓													
3	↓							↓							↓													
4															↓													
5	↓							↓																				
6	↓							↓							↓													

	day																											
Rp.	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84
1	↓							↓							↓													
2	↓							↓							↓													
3	↓							↓							↓													
4								↓																				
5	↓														↓													
6	↓							↓							↓													

投与順序 day1、43、64：1→2→3→4→6 day8、15、29、36、57、71：1→2→3→5→6 day22：1→4