

Weekly パクリタキセル

適応疾患：食道癌

1コースの期間：7週間 上限コース回数：なし

Rp.	薬剤	投与量	投与経路	投与速度、時間
1	生食	250mL	点滴静注	メインルート
2	レスタミンコーワ錠	5錠(50mg)	内服	パクリタキセル投与30分前までに
3	デキサート注	8.25mg	静注	パクリタキセル投与約30分前に 緩徐に静注
	ファモチジン注	20mg		
4	生食	50mL	点滴静注	30分
5	パクリタキセル	100mg/m ²	点滴静注	60分
	生食	250mL		

		day																												
Rp.		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
1	↓								↓							↓							↓							
2	↓								↓							↓							↓							
3	↓								↓							↓							↓							
4	↓								↓							↓							↓							
5	↓								↓							↓							↓							

		day																												
Rp.		29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49								
1	↓								↓																					
2	↓								↓																					
3	↓								↓																					
4	↓								↓																					
5	↓								↓																					

投与順序 1→2→3→4→5