

ザノサー

適応疾患：消化管神経内分泌腫瘍

1コースの期間：6週間 上限コース回数：なし

Rp.	薬剤	投与量	投与経路	投与速度、時間
1	アブレピタントカプセル	day1:125mg day2、3:80mg	内服	抗がん剤投与60～90分前 午前中
2	生食	500mL	点滴静注	メインルート
3	グラニセロン静注	3mg	点滴静注	15分
	デキサート注	9.9mg		
	生食	50mL		
4	グラニセロン静注	3mg	点滴静注	15分
	デキサート注	6.6mg		
	生食	50mL		
5	ザノサー	500mg/m ²	点滴静注	30分
	生食	100mL		
6	生食	250mL	点滴静注	60分

		day																											
Rp.		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
1		↓	↓	↓																									
2		↓	↓	↓	↓	↓																							
3		↓																											
4			↓	↓	↓	↓																							
5		↓	↓	↓	↓	↓																							
6		↓	↓	↓	↓	↓																							

		day													
Rp.		29	30	31	32	33	34	35	36	37	38	39	40	41	42
1															
2															
3															
4															
5															
6															

投与順序 1日目:1→2→3→5→6 2-5日目:2→4→5→6